

Product Description

293-ES cells are derived from the human embryonic kidney HEK 293 cell line. 293-ES cells are adapted for suspension culture in ESF SFM and are available as a frozen vial or suspension culture.

For Research Use Only. Not for use in diagnostic procedures.

Product	Catalog Number	Amount	Storage
293-ES cells adapted in ESF SFM, frozen vial	94-007F	20 million cells per vial	Thaw immediately or LN ₂
293-ES cells adapted in ESF SFM, suspension culture	94-007S	20 million cells in 15 ml media	Culture immediately

Important Information

ESF SFM is a 1X complete, ready to use medium. Do not add L-Glutamine or surfactants such as Pluronic® F-68. Antibiotics are not recommended; however, Penicillin-Streptomycin or Gentamicin may be used when required.

Safety Information

Read the Safety Data Sheets (SDS) and follow the handling instructions. Wear appropriate protective eyewear, clothing and gloves.

Culture Conditions

Media: ESF SFM

Cell Line(s): 293-ES

Culture Type: Suspension or adherent

Recommended Culture Vessels: Shake flasks (vented is recommended) or spinner bottle

Temperature Range: 37°C

Incubator Atmosphere: Humidified, 5% CO₂ atmosphere. Ensure proper gas exchange and minimize exposure of cultures to light.

Receiving Frozen Cells

293-ES cells are frozen in ESF SFM with 10% DMSO. There are 20 x 10⁶ cells per vial. Prepare for thawing cells by placing 20 ml pre-warmed ESF SFM into a 125 ml vented Erlenmeyer shake flask. Thaw frozen cells rapidly by shaking in a 37°C water bath. Thaw vial until a small amount of ice remains. Do not leave vial unattended. Transfer contents of vial to culture flasks using a 1 ml pipette. Do not pour. Incubate overnight at 37°C in a humidified, 5% CO₂ shaking incubator. Determine count and viability and bring volume up to 40 ml total using ESF SFM.

Receiving Suspension Cultures

293-ES cells are packaged in a 15 ml conical filled to the top with ESF SFM. There are 20 x 10⁶ cells per conical. Take extreme care when removing the lid of the conical. Transfer the contents of the conical to a 125 ml vented Erlenmeyer shake flask using a 10 mL pipette. Do not pour. Bring the volume up to 40 ml with pre-warmed ESF SFM.

Suspension Cell Culture

It is recommended to passage the cells three days a week on a Mon/Wed/Fri schedule or twice a week on a Mon/Thurs or Tues/Fri schedule. It is not advised to repeatedly allow the cells to reach maximum densities as the growth kinetics of the culture

may change. Try to keep the maximum cell density to mid-log phase.

Note: It is recommended that a growth curve be determined using the user's standard culturing conditions. This will allow for determination of mid-log phase growth.

1. Determine viable cell count.
2. Seed shake flask between 0.3-0.5 x 10⁶ cell/mL. Use 30-40 mL for a 125 mL Erlenmeyer shake flask, 50-75 mL for 100 mL spinner bottle.
3. Incubate at 37°C in a humidified, 5% CO₂ atmosphere incubator. Rotate shake flask cultures on an orbital shaker platform at 140-150 rpm. Use vented 125 ml Erlenmeyer flasks to allow for gas exchange (recommended) or loosen caps. For spinner cultures, set impeller stirring rate to 85-95 rpm (rpm may vary with impeller design). Loosen side arm caps to allow for gas exchange.
4. Passage when viable cell density reaches 3 x 10⁶ cells/mL. Cultures will grow to densities in excess of 4 x 10⁶ but repeated passage at high densities is not recommended.
5. It is recommended to thaw a new vial of cells every 3 months. Cultures may be maintained for a longer time period but increase the risk of accumulating environmental stresses that can impact the growth and performance characteristics of the culture.

Monolayer Cell Culture

1. Observe cell monolayer using an inverted microscope to ensure confluence. Remove media and any floating cells using a sterile pipette or by aspiration.
2. Add 4 mL (per 25 cm²) ESF SFM to the flask and resuspend the cells by repeatedly pipetting the medium across the monolayer. It may be necessary to aid cell detachment by tapping the side of the flask against a hard surface.
3. Determine the viable cell density of the cell suspension.
4. Inoculate 0.5-1 x 10⁶ cells (per 25 cm²) into new culture flasks containing room temperature ESF SFM (5 mL per 25 cm²).
5. Incubate at 37°C in a humidified, 5% CO₂ atmosphere incubator. Loosen caps or use flasks with vented caps (recommended).








Cryopreservation

- Freezing medium is sterile filtered 90% ESF SFM plus 10% DMSO. 0.15 M trehalose may be added. Store and use at 4°C.
- Prepare the desired quantity of cells, harvesting in mid-log growth with viability >90%.
- Determine the viable cell density and calculate the required volume of freezing medium to give a final cell density between 10-20 x 10⁶ cells/mL.
- Harvest the cells by centrifugation at 1000 rpm for 5 minutes. Resuspend the cells in the pre-determined volume of 4°C freezing medium.
- Dispense 1 mL aliquots of suspension into cryovials.
- Achieve cryopreservation in an automated or manual controlled rate freezing apparatus following standard procedures (1°C decrease per minute).
- Transfer frozen cells to liquid nitrogen, we recommend vapor phase storage at -200 °C to -125 °C.

Related Products

Product	Catalog Number
ESF SFM	98-001
ESF 921	96-001
Adapted CHO Cells	94-008

Legend of Labeling Symbols

Symbol	Interpretation
	Catalog Number
	Lot Number
	<i>Research Use Only</i>
	Manufacturer
	Temperature Limitation
	Date of Manufacture
	Instruction for Use

Important Licensing Information

This product may be covered by one or more Limited Use Label Licenses. By use of this product, you accept the terms and conditions of all applicable Limited Use Label Licenses.

Limited Product Warranty

Expression Systems LLC warrants that this product meets its specifications, as stated in our product brochures and certificates. This warranty lasts from the time we deliver the consumable until either the consumable's shelf life, when the product has been handled and stored in accordance with this IFU.

For technical assistance or documentation, such as Certificates of Analysis or Safety Data Sheets, email support@expressionsystems.com

©2016 Expression Systems LLC. All rights reserved.

Kolliphor® (Pluronic® F-68) is a registered trademark of BASF Corporation.

DISCLAIMER – EXPRESSION SYSTEMS LLC DISCLAIMS ALL WARRANTIES WITH RESPECT TO THIS DOCUMENT, EXPRESSED OR IMPLIED, INCLUDING BUT NOT LIMITED TO THOSE OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR NON-INFRINGEMENT. TO THE EXTENT ALLOWED BY LAW, IN NO EVENT SHALL EXPRESSION SYSTEMS AND/OR ITS AFFILIATE(S) BE LIABLE, WHETHER IN CONTRACT, TORT, WARRANTY, OR UNDER ANY STATUTE OR ON ANY OTHER BASIS FOR SPECIAL, INCIDENTAL, INDIRECT, PUNITIVE, MULTIPLE OR CONSEQUENTIAL DAMAGES IN CONNECTION WITH OR ARISING FROM THIS DOCUMENT, INCLUDING BUT NOT LIMITED TO THE USE THEREOF.

